

Student Name: \_\_\_\_\_ Date of Test: \_\_\_\_\_  
 Hospital Scenario: "You are working in a hospital or clinic when a woman runs through the door, carrying an infant. She shouts, 'Help me! My baby's not breathing.' You have gloves and a pocket mask. You send your coworker to activate the emergency response system and to get the emergency equipment."  
 Prehospital Scenario: "You arrive on the scene for an infant who is not breathing. No bystander CPR has been provided. You approach the scene and ensure that it is safe. Demonstrate what you would do next."

**Assessment and Activation**  
 Checks responsiveness  
 Checks breathing  
 Shouts for help/Activates emergency response system  
 Checks pulse

*Once students shout for help, instructor says, "Here's the barrier device."*

**Cycle 1 of CPR (30:2) \*CPR feedback devices preferred for accuracy**

**Infant Compressions**

- Performs high-quality compressions\*:
  - Placement of 2 fingers or 2 thumbs in the center of the chest, just below the nipple line
  - 30 compressions in no less than 15 and no more than 18 seconds
  - Compresses at least one third the depth of the chest, about 1½ inches (4 cm)
  - Complete recoil after each compression

**Infant Breaths**

- Gives 2 breaths with a barrier device:
  - Each breath given over 1 second
  - Visible chest rise with each breath
- Resumes compressions in less than 10 seconds

**Cycle 2 of CPR (repeats steps in Cycle 1) Only check box if step is successfully performed**

Compressions  Breaths  Resumes compressions in less than 10 seconds

*Rescuer 2 arrives with bag-mask device and begins ventilation while Rescuer 1 continues compressions with 2 thumb—encircling hands technique.*

**Cycle 3 of CPR**

- Powers on AED  Correctly attaches pads  Clears for analysis  Clears to safely deliver a shock
- Safely delivers a shock

**Rescuer 1: Infant Compressions**

- Performs high-quality compressions\*:
  - 15 compressions with 2 thumb—encircling hands technique
  - 15 compressions in no less than 7 and no more than 9 seconds
  - Compress at least one third the depth of the chest, about 1½ inches (4 cm)
- Complete recoil after each compression

**Rescuer 2: Infant Breaths**

*This rescuer is not evaluated.*

**Cycle 4 of CPR**

**Rescuer 2: Infant Compressions**

*This rescuer is not evaluated.*

**Rescuer 1: Infant Breaths**

- Gives 2 breaths with a bag-mask device:
  - Each breath given over 1 second
  - Visible chest rise with each breath
- Resumes compressions in less than 10 seconds

**Instructor Notes**

- Place a check in the box next to each step the student completes successfully.
- If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation).

**Test Results** Check **PASS** or **NR** to indicate pass or needs remediation:  **PASS**  **NR**

Instructor Initials: \_\_\_\_\_ Instructor Number: \_\_\_\_\_ Date: \_\_\_\_\_

©2020 American Heart Association

Student Name: \_\_\_\_\_ Date of Test: \_\_\_\_\_  
 Hospital Scenario: "You are working in a hospital or clinic, and you see a child who has suddenly collapsed in the hallway. You check that the scene is safe and then approach the patient. Demonstrate what you would do next."  
 Prehospital Scenario: "You arrive on the scene for a child who is not breathing. No bystander CPR has been provided. You approach the scene and ensure that it is safe. Demonstrate what you would do next."

**Assessment and Activation**  
 Checks responsiveness  
 Checks breathing  
 Shouts for help/Activates emergency response system/Sends for AED  
 Checks pulse

*Once students shout for help, instructor says, "Here's the barrier device. I'm going to get the AED."*

**Cycle 1 of CPR (30:2) \*CPR feedback devices preferred for accuracy**

**Child Compressions**

- Performs high-quality compressions\*:
  - Hand placement on lower half of sternum
  - 30 compressions in no less than 15 and no more than 18 seconds
  - Compresses at least one third the depth of the chest, about 2 inches (5 cm)
  - Complete recoil after each compression

**Child Breaths**

- Gives 2 breaths with a barrier device:
  - Each breath given over 1 second
  - Visible chest rise with each breath
- Resumes compressions in less than 10 seconds

**Cycle 2 of CPR (repeats steps in Cycle 1) Only check box if step is successfully performed**

Compressions  Breaths  Resumes compressions in less than 10 seconds

*Rescuer 2 says, "Here is the AED. I'll take over compressions, and you use the AED."*

**AED (follows prompts of AED)**

- Powers on AED  Correctly attaches pads  Clears for analysis  Clears to safely deliver a shock
- Safely delivers a shock

**Resumes Compressions**

- Ensures compressions are resumed immediately after shock delivery
- Student directs instructor to resume compressions or
- Student resumes compressions

**STOP TEST**

**Instructor Notes**

- Place a check in the box next to each step the student completes successfully.
- If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation).

**Test Results** Check **PASS** or **NR** to indicate pass or needs remediation:  **PASS**  **NR**

Instructor Initials: \_\_\_\_\_ Instructor Number: \_\_\_\_\_ Date: \_\_\_\_\_

©2020 American Heart Association